

Math 20200

Calculus II

Lesson 8

Hyperbolic Functions

Dr. A. Marchese, The City College of New York



Table of Contents:

1. Definition of $\cosh(t)$ and $\sinh(t)$	00:46	p.2
2. The other hyperbolic functions	05:00	p.4
3. Derivatives of hyperbolic functions	06:50	p.5
4. Catenary curves	16:30	p.10

Hyperbolic Functions

Consider $\frac{e^x - e^{-x}}{2}$ and $\frac{e^x + e^{-x}}{2}$.

These functions have the same relationship to the hyperbola $x^2 - y^2 = 1$ as $y = \cos x$ and $y = \sin x$ have with the unit circle $x^2 + y^2 = 1$.



$$\therefore \cos^2 t + \sin^2 t = 1$$

call $\frac{e^t + e^{-t}}{2} = \cosh t$

$$\frac{e^t - e^{-t}}{2} = \sinh t$$

hyperbolic cosine +
hyperbolic sine

$$\therefore \cosh^2 t - \sinh^2 t = 1 .$$

Ex. Evaluate $\sinh(3)$

$$\sinh(3) = \frac{e^3 - e^{-3}}{2}$$

Ex. Evaluate $\cosh(\ln 3)$

$$\cosh(\ln 3) = \frac{e^{\ln 3} + e^{-\ln 3}}{2} =$$



Work on this problem
on your own

$$\begin{aligned} e^{-\ln 3} &= e^{\ln(3^{-1})} \\ &= e^{\ln(\frac{1}{3})} \\ &= \frac{1}{3} \end{aligned}$$

$$= \frac{3 + (e^{\ln 3})^{-1}}{2} = \frac{3 + 3^{-1}}{2} = \frac{3 + \frac{1}{3}}{2}$$

$$\frac{1}{2}\sqrt{30} = \frac{\sqrt{10}}{6} = \boxed{\frac{5}{3}}$$

$$\text{Ex. } \sinh(0) = \frac{e^0 - e^{-0}}{2} = \frac{1-1}{2} = 0$$

$$\cosh(0) = \frac{e^0 + e^{-0}}{2} = \frac{1+1}{2} = \frac{2}{2} = 1.$$

Ex. prove $\sinh(-x) = -\sinh(x)$.

$$\begin{aligned}
 \sinh(-x) &= \frac{e^{-x} - e^{-(-x)}}{2} \\
 &= \frac{e^{-x} - e^x}{2} = -\left(\frac{e^x - e^{-x}}{2}\right) \\
 &= -\sinh x
 \end{aligned}$$

$$\text{So we have : } \sinh x = \frac{e^x - e^{-x}}{2}, \cosh x = \frac{e^x + e^{-x}}{2}$$

and we define $\tanh x = \frac{\sinh x}{\cosh x}$, $\coth x = \frac{\cosh x}{\sinh x}$

$$\operatorname{Sech} x = \frac{1}{\cosh x} \quad \text{and} \quad \operatorname{Csch} x = \frac{1}{\sinh x}$$

Ex. If $\sinh(x) = \frac{3}{4}$ find the values of $\cosh(x)$ and $\tanh(x)$.

To find $\cosh x$, use $\cosh^2 x - \sinh^2 x = 1$

$$\cosh^2 x - \left(\frac{3}{4}\right)^2 = 1$$

$$\cosh^2 x - \frac{9}{16} = 1$$

$$\cosh^2 x = 1 + \frac{9}{16} = \frac{25}{16}$$

$$\cosh x = \pm \sqrt{\frac{25}{16}}$$

$$= \pm \frac{5}{4}$$

\cosh must be positive

$$\frac{e^x + e^{-x}}{2}$$

then $\tanh x = \frac{\sinh x}{\cosh x} = \frac{\frac{3}{4}}{\frac{5}{4}} = \boxed{\frac{3}{5}}$.

Derivatives of Hyperbolic Functions :

$$\begin{aligned} \frac{d}{dx}(\sinh x) &= \frac{d}{dx}\left(\frac{e^x - e^{-x}}{2}\right) = \frac{d}{dx}\left(\frac{1}{2}(e^x - e^{-x})\right) \\ &= \frac{1}{2}\left(e^x - \cancel{e^{-x}}(-1)\right) = \frac{1}{2}(e^x + e^{-x}) = \cosh x. \end{aligned}$$

Similarly, $\frac{d}{dx}(\cosh x) = \sinh x$ (exercise)


Ex. $f(t) = \ln(\sinh t)$

$$f'(t) = \frac{1}{\sinh t} \cdot \frac{d}{dt}(\sinh t) = \frac{\cosh t}{\sinh t}$$
$$= \coth(t).$$

Ex. $y = \sinh(\cosh x)$

$$y' = \cosh(\cosh x) \cdot \frac{d}{dx}(\cosh x) \quad \text{chain rule}$$
$$= \cosh(\cosh x) \cdot \sinh x$$
$$= \sinh x \cdot \cosh(\cosh x).$$

Ex. $\int \frac{\sinh x}{1 + \cosh^2 x} dx$



Work on this problem
on your own

$$\int \frac{\sinh x}{1 + \cosh^2 x} dx = \int \frac{\sinh x}{1 + (\cosh x)^2} dx$$

$u = \cosh x$
 $du = \sinh x dx$

$$= \int \frac{1}{1 + u^2} du = \arctan u + C$$

$$= \arctan(\cosh x) + C.$$

Since $\frac{d}{dx}(\sinh x) = \cosh x > 0 \quad \forall x$, we know

$\sinh x$ is one to one and hence has an inverse.

Ex. Prove that $\sinh^{-1} x = \ln(x + \sqrt{x^2 + 1})$

2 ways we can do this:

1) start with $y = \sinh x$ and find the inverse function

2) let $f(x) = \sinh x$ and $g(x) = \ln(x + \sqrt{x^2 + 1})$

and show $f(g(x)) = x$ and $g(f(x)) = x \quad \forall x \in \mathbb{R}$

1) $y = \sinh x = \frac{e^x - e^{-x}}{2}$ solve for x

$$2y = e^x - e^{-x} \quad \text{mult. by } e^x$$

$$2ye^x = e^{2x} - 1$$

$$0 = e^{2x} - 2ye^x - 1 \quad \text{treat as a quadratic}$$

$$0 = (e^x)^2 - 2y e^x - 1 \quad a=1 \quad b=-2y \quad c=-1$$

$$e^x = \frac{2y \pm \sqrt{4y^2 + 4}}{2} = y \pm \sqrt{y^2 + 1}$$

notice e^x must be positive, so we

can't have $y - \sqrt{y^2 + 1}$ ($\sqrt{y^2 + 1} > y$)

$$\therefore e^x = y + \sqrt{y^2 + 1}$$

$$x = \ln(y + \sqrt{y^2 + 1}) \quad \text{now switch } x \leftrightarrow y$$

$$y = \ln(x + \sqrt{x^2 + 1}) = \sinh^{-1} x.$$

$$2) \quad f(x) = \sinh x = \frac{e^x - e^{-x}}{2} \quad g(x) = \ln(x + \sqrt{x^2 + 1})$$

$$f(g(x)) = \frac{e^{\ln(x + \sqrt{x^2 + 1})} - e^{-\ln(x + \sqrt{x^2 + 1})}}{2} =$$

$$= \frac{(x + \sqrt{x^2+1}) - e^{\ln(\frac{1}{x+\sqrt{x^2+1}})}}{2}$$

$$= \frac{x + \sqrt{x^2+1} - \frac{1}{x+\sqrt{x^2+1}}}{2} \cdot \frac{x + \sqrt{x^2+1}}{x + \sqrt{x^2+1}}$$

$$= \frac{x^2 + 2x\sqrt{x^2+1} + x^2+1 - 1}{2(x+\sqrt{x^2+1})}$$

$$= \frac{2x^2 + 2x\sqrt{x^2+1}}{2(x+\sqrt{x^2+1})} = \frac{2x(x + \sqrt{x^2+1})}{2(x+\sqrt{x^2+1})} = x$$

$$g(f(x)) = \ln\left(\frac{e^x - e^{-x}}{2} + \sqrt{\left(\frac{e^x - e^{-x}}{2}\right)^2 + 1}\right)$$

$$= \ln\left(\frac{e^x - e^{-x}}{2} + \sqrt{\frac{e^{2x} - 2 + e^{-2x}}{4} + 1}\right) - \frac{2}{4} + \frac{4}{4} = +\frac{2}{4}$$

$$= \ln\left(\frac{e^x - e^{-x}}{2} + \sqrt{\frac{e^{2x} + 2 + e^{-2x}}{4}}\right)$$

$$= \ln\left(\frac{e^x - e^{-x}}{2} + \sqrt{\left(\frac{e^x + e^{-x}}{2}\right)^2}\right)$$


$$\begin{aligned}
 &= \ln \left(\frac{e^x - e^{-x}}{2} + \frac{e^x + e^{-x}}{2} \right) \\
 &= \ln \left(\frac{2e^x}{2} \right) = \ln(e^x) = x .
 \end{aligned}$$

Catenary curves :

A flexible cable always hangs in the shape of a catenary curve

$$y = c + a \cosh\left(\frac{x}{a}\right) \quad c, a \text{ constants}$$

$$a > 0$$



many applications, including architecture :

St. Louis Gateway Arch,

works of Antoni Gaudi ←

google :

antonio gaudi catenary