

Decoding the rules:  $\Delta x = \frac{b-a}{n}$      $x_i = a + i\Delta x$      $\bar{x}_i = \frac{1}{2}(x_{i-1} + x_i)$

Midpoint Rule:  $\int_a^b f(x) dx \approx M_n = \frac{\Delta x}{1} [f(\bar{x}_1) + f(\bar{x}_2) + f(\bar{x}_3) + \dots + f(\bar{x}_{n-1}) + f(\bar{x}_n)]$

Trapezoidal Rule:  $\int_a^b f(x) dx \approx T_n = \frac{\Delta x}{2} [f(x_0) + 2f(x_1) + 2f(x_2) + 2f(x_3) + \dots + 2f(x_{n-1}) + f(x_n)]$

Simpson's Rule:  $\int_a^b f(x) dx \approx S_n = \frac{\Delta x}{3} [f(x_0) + 4f(x_1) + 2f(x_2) + 4f(x_3) + \dots + 2f(x_{n-2}) + 4f(x_{n-1}) + f(x_n)]$

The value of  $n$  must be even for Simpson's rule to work.

Error Bounds: For  $a \leq x \leq b$

Midpoint error:  $|E_M| \leq \frac{K(b-a)^3}{24n^2}$      $|f''(x)| \leq K$

Trapezoidal error:  $|E_T| \leq \frac{K(b-a)^3}{12n^2}$      $|f''(x)| \leq K$

Simpson error:  $|E_S| \leq \frac{K(b-a)^5}{180n^4}$      $|f^{(4)}(x)| \leq K$

Additional examples:

8)  $\int_0^2 \frac{1}{1+x^6} dx$      $n = 8$

$\Delta x = \frac{(2)-(0)}{(8)} = \frac{2}{8} = \frac{1}{4}$      $\frac{\Delta x}{2} = \frac{1}{8}$

$\frac{0}{8}$	$\frac{1}{8}$		$\frac{3}{8}$		$\frac{5}{8}$		$\frac{7}{8}$		$\frac{9}{8}$		$\frac{11}{8}$		$\frac{13}{8}$		$\frac{15}{8}$	
$x_0$	$\bar{x}_1$	$x_1$	$\bar{x}_2$	$x_2$	$\bar{x}_3$	$x_3$	$\bar{x}_4$	$x_4$	$\bar{x}_5$	$x_5$	$\bar{x}_6$	$x_6$	$\bar{x}_7$	$x_7$	$\bar{x}_8$	$x_8$
$0 = \frac{0}{4}$		$\frac{1}{4}$		$\frac{2}{4}$		$\frac{3}{4}$		$\frac{4}{4}$		$\frac{5}{4}$		$\frac{6}{4}$		$\frac{7}{4}$		$\frac{8}{4}$


$$M_8 = \frac{\left(\frac{1}{4}\right)}{1} \left[ \frac{1}{1+\left(\frac{1}{8}\right)^6} + \frac{1}{1+\left(\frac{3}{8}\right)^6} + \frac{1}{1+\left(\frac{5}{8}\right)^6} + \frac{1}{1+\left(\frac{7}{8}\right)^6} + \frac{1}{1+\left(\frac{9}{8}\right)^6} + \frac{1}{1+\left(\frac{11}{8}\right)^6} + \frac{1}{1+\left(\frac{13}{8}\right)^6} + \frac{1}{1+\left(\frac{15}{8}\right)^6} \right]$$

$$T_8 = \frac{\left(\frac{1}{4}\right)}{2} \left[ \frac{1}{1+\left(\frac{1}{4}\right)^6} + 2\left(\frac{1}{1+\left(\frac{2}{4}\right)^6}\right) + 2\left(\frac{1}{1+\left(\frac{3}{4}\right)^6}\right) + 2\left(\frac{1}{1+\left(\frac{4}{4}\right)^6}\right) \right. \\ \left. + 2\left(\frac{1}{1+\left(\frac{5}{4}\right)^6}\right) + 2\left(\frac{1}{1+\left(\frac{6}{4}\right)^6}\right) + 2\left(\frac{1}{1+\left(\frac{7}{4}\right)^6}\right) + \frac{1}{1+\left(\frac{8}{4}\right)^6} \right]$$

$$S_8 = \frac{\left(\frac{1}{4}\right)}{3} \left[ \frac{1}{1+\left(\frac{1}{4}\right)^6} + 4\left(\frac{1}{1+\left(\frac{2}{4}\right)^6}\right) + 2\left(\frac{1}{1+\left(\frac{3}{4}\right)^6}\right) + 4\left(\frac{1}{1+\left(\frac{4}{4}\right)^6}\right) \right. \\ \left. + 2\left(\frac{1}{1+\left(\frac{5}{4}\right)^6}\right) + 4\left(\frac{1}{1+\left(\frac{6}{4}\right)^6}\right) + 2\left(\frac{1}{1+\left(\frac{7}{4}\right)^6}\right) + \frac{1}{1+\left(\frac{8}{4}\right)^6} \right]$$

12)  $\int_0^1 \sin(x^3) dx \quad n=10$

$$\Delta x = \frac{(1)-(0)}{(10)} = \frac{1}{10} = \frac{2}{20} \quad \frac{\Delta x}{2} = \frac{1}{20}$$




$$M_{10} = \frac{\left(\frac{1}{10}\right)}{1} \left[ \sin\left(\frac{1}{20}\right) + \sin\left(\frac{3}{20}\right) + \sin\left(\frac{5}{20}\right) + \sin\left(\frac{7}{20}\right) + \sin\left(\frac{9}{20}\right) \right. \\ \left. + \sin\left(\frac{11}{20}\right) + \sin\left(\frac{13}{20}\right) + \sin\left(\frac{15}{20}\right) + \sin\left(\frac{17}{20}\right) + \sin\left(\frac{19}{20}\right) \right]$$

$$T_{10} = \frac{\left(\frac{1}{10}\right)}{2} \left[ \sin\left(\frac{0}{10}\right) + 2\left(\sin\left(\frac{1}{10}\right)\right) + 2\left(\sin\left(\frac{2}{10}\right)\right) + 2\left(\sin\left(\frac{3}{10}\right)\right) + 2\left(\sin\left(\frac{4}{10}\right)\right) + 2\left(\sin\left(\frac{5}{10}\right)\right) \right. \\ \left. + 2\left(\sin\left(\frac{6}{10}\right)\right) + 2\left(\sin\left(\frac{7}{10}\right)\right) + 2\left(\sin\left(\frac{8}{10}\right)\right) + 2\left(\sin\left(\frac{9}{10}\right)\right) + \sin\left(\frac{10}{10}\right) \right]$$

$$S_{10} = \frac{\left(\frac{1}{10}\right)}{2} \left[ \sin\left(\frac{0}{10}\right) + 4\left(\sin\left(\frac{1}{10}\right)\right) + 2\left(\sin\left(\frac{2}{10}\right)\right) + 4\left(\sin\left(\frac{3}{10}\right)\right) + 2\left(\sin\left(\frac{4}{10}\right)\right) + 4\left(\sin\left(\frac{5}{10}\right)\right) \right. \\ \left. + 2\left(\sin\left(\frac{6}{10}\right)\right) + 4\left(\sin\left(\frac{7}{10}\right)\right) + 2\left(\sin\left(\frac{8}{10}\right)\right) + 4\left(\sin\left(\frac{9}{10}\right)\right) + \sin\left(\frac{10}{10}\right) \right]$$

14)  $\int_4^6 \ln(x^3 + 2) dx \quad n = 10$

$$\Delta x = \frac{(6)-(4)}{(10)} = \frac{2}{10} = \frac{1}{5} \quad \frac{\Delta x}{2} = \frac{1}{10}$$



$$M_{10} = \frac{\left(\frac{1}{5}\right)}{1} \left[ \ln\left(\left(\frac{41}{10}\right)^3 + 2\right) + \ln\left(\left(\frac{43}{10}\right)^3 + 2\right) + \ln\left(\left(\frac{45}{10}\right)^3 + 2\right) + \ln\left(\left(\frac{47}{10}\right)^3 + 2\right) + \ln\left(\left(\frac{49}{10}\right)^3 + 2\right) \right. \\ \left. + \ln\left(\left(\frac{51}{10}\right)^3 + 2\right) + \ln\left(\left(\frac{53}{10}\right)^3 + 2\right) + \ln\left(\left(\frac{55}{10}\right)^3 + 2\right) + \ln\left(\left(\frac{57}{10}\right)^3 + 2\right) + \ln\left(\left(\frac{59}{10}\right)^3 + 2\right) \right]$$



$$M_{10} = \frac{\left(\frac{1}{10}\right)}{1} \left[ \begin{aligned} &\frac{\sqrt{\left(\frac{1}{20}\right)}}{e^{\left(\frac{1}{20}\right)}} + \frac{\sqrt{\left(\frac{3}{20}\right)}}{e^{\left(\frac{3}{20}\right)}} + \frac{\sqrt{\left(\frac{5}{20}\right)}}{e^{\left(\frac{5}{20}\right)}} + \frac{\sqrt{\left(\frac{7}{20}\right)}}{e^{\left(\frac{7}{20}\right)}} + \frac{\sqrt{\left(\frac{9}{20}\right)}}{e^{\left(\frac{9}{20}\right)}} \\ &+ \frac{\sqrt{\left(\frac{11}{20}\right)}}{e^{\left(\frac{11}{20}\right)}} + \frac{\sqrt{\left(\frac{13}{20}\right)}}{e^{\left(\frac{13}{20}\right)}} + \frac{\sqrt{\left(\frac{15}{20}\right)}}{e^{\left(\frac{15}{20}\right)}} + \frac{\sqrt{\left(\frac{17}{20}\right)}}{e^{\left(\frac{17}{20}\right)}} + \frac{\sqrt{\left(\frac{19}{20}\right)}}{e^{\left(\frac{19}{20}\right)}} \end{aligned} \right]$$

$$T_{10} = \frac{\left(\frac{1}{10}\right)}{2} \left[ \begin{aligned} &\frac{\sqrt{\left(\frac{0}{10}\right)}}{e^{\left(\frac{0}{10}\right)}} + 2 \left( \frac{\sqrt{\left(\frac{1}{10}\right)}}{e^{\left(\frac{1}{10}\right)}} \right) + 2 \left( \frac{\sqrt{\left(\frac{2}{10}\right)}}{e^{\left(\frac{2}{10}\right)}} \right) + 2 \left( \frac{\sqrt{\left(\frac{3}{10}\right)}}{e^{\left(\frac{3}{10}\right)}} \right) + 2 \left( \frac{\sqrt{\left(\frac{4}{10}\right)}}{e^{\left(\frac{4}{10}\right)}} \right) + 2 \left( \frac{\sqrt{\left(\frac{5}{10}\right)}}{e^{\left(\frac{5}{10}\right)}} \right) \\ &+ 2 \left( \frac{\sqrt{\left(\frac{6}{10}\right)}}{e^{\left(\frac{6}{10}\right)}} \right) + 2 \left( \frac{\sqrt{\left(\frac{7}{10}\right)}}{e^{\left(\frac{7}{10}\right)}} \right) + 2 \left( \frac{\sqrt{\left(\frac{8}{10}\right)}}{e^{\left(\frac{8}{10}\right)}} \right) + 2 \left( \frac{\sqrt{\left(\frac{9}{10}\right)}}{e^{\left(\frac{9}{10}\right)}} \right) + \frac{\sqrt{\left(\frac{10}{10}\right)}}{e^{\left(\frac{10}{10}\right)}} \end{aligned} \right]$$

$$S_{10} = \frac{\left(\frac{1}{10}\right)}{2} \left[ \begin{aligned} &\frac{\sqrt{\left(\frac{0}{10}\right)}}{e^{\left(\frac{0}{10}\right)}} + 4 \left( \frac{\sqrt{\left(\frac{1}{10}\right)}}{e^{\left(\frac{1}{10}\right)}} \right) + 2 \left( \frac{\sqrt{\left(\frac{2}{10}\right)}}{e^{\left(\frac{2}{10}\right)}} \right) + 4 \left( \frac{\sqrt{\left(\frac{3}{10}\right)}}{e^{\left(\frac{3}{10}\right)}} \right) + 2 \left( \frac{\sqrt{\left(\frac{4}{10}\right)}}{e^{\left(\frac{4}{10}\right)}} \right) + 4 \left( \frac{\sqrt{\left(\frac{5}{10}\right)}}{e^{\left(\frac{5}{10}\right)}} \right) \\ &+ 2 \left( \frac{\sqrt{\left(\frac{6}{10}\right)}}{e^{\left(\frac{6}{10}\right)}} \right) + 4 \left( \frac{\sqrt{\left(\frac{7}{10}\right)}}{e^{\left(\frac{7}{10}\right)}} \right) + 2 \left( \frac{\sqrt{\left(\frac{8}{10}\right)}}{e^{\left(\frac{8}{10}\right)}} \right) + 4 \left( \frac{\sqrt{\left(\frac{9}{10}\right)}}{e^{\left(\frac{9}{10}\right)}} \right) + \frac{\sqrt{\left(\frac{10}{10}\right)}}{e^{\left(\frac{10}{10}\right)}} \end{aligned} \right]$$

20)  $\int_0^1 e^{x^2} dx \quad |E_S| \leq 0.00001$

$y = e^{x^2} \quad f^{(4)}(x) = \frac{d^4 y}{dx^4} = e^{x^2} (12 + 48x^2 + 16x^4) \quad |E_S| \leq \frac{K(b-a)^5}{180n^4} \quad |f^{(4)}(x)| \leq K$

$|f^{(4)}(x)| \leq K = |e^{(1)^2} (12 + 48(1)^2 + 16(1)^4)| = 76e$

$$0.00001 \leq \frac{(76e)((1)-(0))^5}{180n^4} \Rightarrow n^4 = \frac{(76e)(100000)}{180} \Rightarrow n = \sqrt[4]{\frac{380000e}{9}}$$

$$n = \sqrt[4]{\frac{380000e}{9}} \Rightarrow n = 18.40597773$$

$$n \approx 20$$