

$$6) \quad r = \sqrt{x^2 + y^2} \quad \theta = \tan^{-1} \frac{y}{x}$$

Part a)  $(3\sqrt{3}, 3)$

$$r = \sqrt{(3\sqrt{3})^2 + (3)^2} = \sqrt{3^2(3+1)} = 3\sqrt{4} = 6$$

$$(i) \quad \theta = \tan^{-1} \left( \frac{3\sqrt{3}}{3} \right) = \tan^{-1} (\sqrt{3}) = \frac{\pi}{6}$$

$$(6, \frac{\pi}{6})$$

$$(ii) \text{ for } r < 0, r = -6 \text{ and } \theta = \frac{\pi}{6} + \pi = \frac{7\pi}{6} \text{ thus } (-6, \frac{7\pi}{6})$$


Part b)  $(1, -2)$

$$r = \sqrt{(1)^2 + (-2)^2} = \sqrt{1+4} = \sqrt{5}$$

$$(i) \quad \theta = \tan^{-1} \left( \frac{-2}{1} \right) = \tan^{-1} (-2)$$

but  $\theta$  is in QIV; therefore  $\theta = 2\pi - \tan^{-1} 2$   
 $(\sqrt{5}, 2\pi - \tan^{-1} 2)$

$$(ii) \text{ for } r < 0, r = -\sqrt{5} \text{ and } \theta = (2\pi - \tan^{-1} 2) - \pi = \pi - \tan^{-1} 2 \text{ thus } (-\sqrt{5}, \pi - \tan^{-1} 2)$$



$$14) \quad \pi = \frac{\pi}{3}$$

$$\frac{y}{x} = \tan \theta = \tan \frac{\pi}{3} = \sqrt{3} \Rightarrow \frac{y}{x} = \sqrt{3} \Rightarrow y = \sqrt{3}x$$

Since we don't have any restriction for  $r$ , this expression  $y = \sqrt{3}x$  with domain  $-\infty < x < \infty$ .

$$16) \quad r = \tan \theta \sec \theta$$

$$r = \tan \theta \sec \theta$$

$$x = r \cos \theta \quad (r \cos \theta)^2 = r \sin \theta$$

$$y = r \sin \theta \quad r = \frac{\sin \theta}{\cos^2 \theta} \Rightarrow (x)^2 = y$$

$$r \cos^2 \theta = \sin \theta \quad y = x^2$$

$$18) \quad 4y^2 = x$$

$$x = r \cos \theta \quad 4y^2 = x \quad \frac{r^2}{r} = \frac{\cos \theta}{4 \sin^2 \theta}$$

$$y = r \sin \theta \quad 4(r \sin \theta)^2 = r \cos \theta \Rightarrow r = \frac{1}{4} \left( \frac{1}{\sin \theta} \right) \left( \frac{\cos \theta}{\sin \theta} \right) = \frac{1}{4} \csc \theta \cot \theta$$

$$20) \quad xy = 4$$

$$x = r \cos \theta \quad xy = 4$$

$$y = r \sin \theta \quad (r \cos \theta)(r \sin \theta) = 4 \Rightarrow r^2 \left( \frac{1}{2} \sin(2\theta) \right) = 4 \quad r^2 = \frac{8}{\sin(2\theta)} = 8 \csc(2\theta)$$

$$r^2 \sin \theta \cos \theta = 4$$



22) a)  $r = 5$  center =  $(2, 3)$

Cartesian is easier:  $(x - 2)^2 + (y - 3)^2 = (5)^2$



b)  $r = 4$  center =  $(0, 0)$

Polar is easier  $r = 4$  instead of Cartesian  $x^2 + y^2 = (4)^2$



28)  $r = \ln \theta$   $\theta \geq 1$





32)  $r = 3 \cos 6\theta$



34)  $r = 2 + \sin \theta$





36)  $r^2 = \cos 4\theta$





Each loop of the Polar plot to the right should have been drawn to the origin.

38)  $r^2\theta = 1$



40)  $r = 3 + 4 \cos \theta$



48)  $r = 2 - \sin \theta \quad \theta = \frac{\pi}{3}$

$$x = r \cos \theta = (2 - \sin \theta) \cos \theta = 2 \cos \theta - \sin \theta \cos \theta = 2 \cos \theta - \frac{1}{2} \sin(2\theta)$$

$$y = r \sin \theta = (2 - \sin \theta) \sin \theta = 2 \sin \theta - \sin^2 \theta$$

$$\frac{dx}{d\theta} = -2 \sin \theta - \frac{1}{2} \cos(2\theta)(2) = -2 \sin \theta - \cos(2\theta) \quad \frac{dy}{d\theta} = 2 \cos \theta - 2 \sin \theta \cos \theta = 2 \cos \theta - \sin(2\theta)$$

$$\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{2 \cos \left(\frac{\pi}{3}\right) - \sin \left(2 \left(\frac{\pi}{3}\right)\right)}{-2 \sin \left(\frac{\pi}{3}\right) - \cos \left(2 \left(\frac{\pi}{3}\right)\right)} = \frac{2 \left(\frac{1}{2}\right) - \left(\frac{\sqrt{3}}{2}\right)}{-2 \left(\frac{\sqrt{3}}{2}\right) - \left(-\frac{1}{2}\right)} = \frac{\frac{2-\sqrt{3}}{2}}{\frac{1-2\sqrt{3}}{2}} = \frac{2-\sqrt{3}}{1-2\sqrt{3}}$$

50)  $r = \cos \left(\frac{\theta}{3}\right) \quad \theta = \pi$

$$x = r \cos \theta = \left(\cos \left(\frac{\theta}{3}\right)\right) \cos \theta = \cos \left(\frac{\theta}{3}\right) \cos \theta \quad y = r \sin \theta = \left(\cos \left(\frac{\theta}{3}\right)\right) \sin \theta = \cos \left(\frac{\theta}{3}\right) \sin \theta$$

$$\frac{dx}{d\theta} = \left[ \frac{-1}{3} \sin \left(\frac{\theta}{3}\right) \right] (\cos \theta) + \left( \cos \left(\frac{\theta}{3}\right) \right) [-\sin \theta] \quad \frac{dy}{d\theta} = \left[ \frac{-1}{3} \sin \left(\frac{\theta}{3}\right) \right] (\sin \theta) + \left( \cos \left(\frac{\theta}{3}\right) \right) [\cos \theta]$$

$$\frac{dy}{dx} = \frac{\frac{-1}{3} \sin \left(\frac{\theta}{3}\right) (\sin \theta) + \left( \cos \left(\frac{\theta}{3}\right) \right) [\cos \theta]}{\frac{-1}{3} \sin \left(\frac{\theta}{3}\right) (\cos \theta) + \left( \cos \left(\frac{\theta}{3}\right) \right) [-\sin \theta]}$$

$$m = \frac{\left[ \frac{-1}{3} \sin \left(\frac{(\pi)}{3}\right) \right] (\sin(\pi)) + \left( \cos \left(\frac{(\pi)}{3}\right) \right) [\cos(\pi)]}{\left[ \frac{-1}{3} \sin \left(\frac{(\pi)}{3}\right) \right] (\cos(\pi)) + \left( \cos \left(\frac{(\pi)}{3}\right) \right) [-\sin(\pi)]} = \frac{\left[ \frac{-1}{3} \left(\frac{\sqrt{3}}{2}\right) \right] (0) + \left(\frac{1}{2}\right)[1]}{\left[ \frac{-1}{3} \left(\frac{\sqrt{3}}{2}\right) \right] (1) + \left(\frac{1}{2}\right)[0]} = \frac{-\frac{1}{2}}{\frac{\sqrt{3}}{3(2)}} = \frac{-3}{\sqrt{3}} = -\sqrt{3}$$