

# SAMPLE 212 FINAL EXAM SOLUTIONS

1. (a) Evaluate:  $\int 5^x \tan(5^x) dx$
- (b) Find  $\frac{d}{dx} \left[ \sqrt{\log_3 x} \right]$
- (c) Does the graph of  $x + x^3 + y^2 + y^4 + z^4 + z^5 = 1$  have symmetry about
- (i) the  $x$ -axis? (ii) the  $xz$ -plane? (iii) the origin?
- (d) What is the value of  $\int_{-\sqrt{\pi}}^{\sqrt{\pi}} \sin(x^3) dx$ . Explain why.

—————SOLUTION—————

(a) Substitute  $u = 5^x$ ;  $du = (\ln 5)5^x dx$ :

$$\int 5^x \tan(5^x) dx = \int \tan u \left( \frac{1}{\ln 5} du \right) = \frac{\ln |\sec(5^x)|}{\ln 5} + C$$

$$\begin{aligned} \text{(b)} \quad \frac{d}{dx} \left[ \sqrt{\log_3 x} \right] &= \frac{1}{2\sqrt{\log_3 x}} \cdot \left( \frac{\ln x}{\ln 3} \right)' \\ &= \frac{1}{2\sqrt{\log_3 x}} \frac{1}{\ln 3} \frac{1}{x} = \frac{1}{2x(\ln 3)\sqrt{\log_3 x}} \\ \text{OR} \quad &= \frac{1}{2\sqrt{(\ln x)/\ln 3}} \frac{1}{\ln 3} \frac{1}{x} = \frac{1}{2x(\sqrt{\ln 3})\sqrt{\ln x}} \end{aligned}$$

(c) (i) Symmetry about the  $x$ -axis: replace  $y$  and  $z$  by their negatives; this is not an equivalent equation, so the answer is No.

(ii)  $xz$ -plane: Replace  $y$  by  $-y$ . Yes.

(iii) origin: Replace all variables by their negative: No

(d) This is an integral of an odd function over symmetric limits. The value is 0.

2. (a) Let the function  $y(x)$  be the solution to the differential equation  $y' = \frac{12x^3 + 12x^2}{y^2 e^{y^3}}$  for which  $y(1) = 0$ . Find  $y(x)$  explicitly as a function of  $x$ .

(b) Find  $\int 4x[\sec(x^2 + 2)] dx$ .

—————SOLUTION—————

(a) Substitute  $u = y^3$ ;  $du = 3y^2 dy$ , let  $C = 3C'$ , and substitute  $x = 1$ :

$$\begin{aligned}\frac{dy}{dx} &= \frac{12x^3 + 12x^2}{y^2 e^{y^3}} \\ \int y^2 e^{y^3} dy &= \int 12x^3 + 12x^2 dx \\ \int e^u \left(\frac{1}{3} du\right) &= \frac{1}{3} e^{y^3} = 3x^4 + 4x^3 + C' \\ y^3 &= \ln(9x^4 + 12x^3 + C) \\ y &= \sqrt[3]{\ln(9x^4 + 12x^3 + C)} \\ 0 &= y(1) = \sqrt[3]{\ln(21 + C)} \\ \ln(21 + C) &= 0; 21 + C = 1 \\ y &= \sqrt[3]{\ln(9x^4 + 12x^3 - 20)}\end{aligned}$$

(b) Let  $u = x^2 + 2$ ;

$$\begin{aligned}\int 4x[\sec(x^2 + 2)] dx &= \int \sec u(2 du) \\ &= 2 \ln |\sec(x^2 + 2) + \tan(x^2 + 2)| + C\end{aligned}$$

3. (a) A package initially with a temperature of  $150^\circ \text{ F}$  is placed in a room maintained at  $70^\circ \text{ F}$ . Assume the temperature difference between package and the room  $t$  hours after the package was initially placed,  $(\Delta T)(t) = T(t) - 70$ , where  $T(t)$  is the temperature of the package  $t$  hours after being placed in the room, satisfies an exponential decay law. The temperature of the package 2 hours after being placed in the room is  $86^\circ \text{ F}$ .

(i) Find the function  $(\Delta T)(t)$ .

(ii) Find the temperature, expressed as a rational number (i.e., a quotient of integers), of the package after 4 hours.

(iii) How long does it take for the package to reach  $71^\circ$ ?

(b) Evaluate:  $\int \sec(3x + 4) dx$

—————SOLUTION—————

$$\begin{aligned} \text{(a,i)} \quad (\Delta T)(t) &= (\Delta T)_0 \left( \frac{(\Delta T)(2)}{(\Delta T)(0)} \right)^{t/(2-0)} \\ &= 80 \left( \frac{16}{80} \right)^{t/2} = 80 \left( \frac{1}{5} \right)^{t/2} = 80e^{-\frac{\ln 5}{2}t} \end{aligned}$$

$$\text{(ii) Using base } \frac{1}{5} \text{ (easiest): } (\Delta T)(4) = 80 \left( \frac{1}{5} \right)^2 = \frac{16}{5}.$$

$$\text{With } e: (\Delta T)(4) = 80e^{-2 \ln 5} = 80e^{\ln(1/25)} = \frac{80}{25} = \frac{16}{5}.$$

$$T(4) = 70 + \frac{16}{5} = 73.2$$

$$\text{(iii)} \quad 1 = (\Delta T)(t) = 80 \left( \frac{1}{5} \right)^{t/2}$$

$$0 = \ln 80 + \frac{t}{2} \ln(1/5)$$

$$t = \frac{-2 \ln 80}{\ln 1/5} = \frac{2 \ln 80}{\ln 5}$$

$$\text{(b) Let } u = 3x + 4; \int \sec(3x + 4) dx = \frac{1}{3} \int \sec u du =$$

$$\frac{1}{3} \ln |\sec(3x + 4) + \tan(3x + 4)| + C$$

4. (a) Evaluate:  $\int_0^1 \arctan x \, dx$

(b) Evaluate:  $\int_1^2 \frac{x^2 - 2x - 4}{x^3 + 2x^2} \, dx$

—————SOLUTION—————

(a) Integrate  $1 \cdot \arctan x$  by parts, with 1 being the factor integrated, and substitute  $u = 1 + x^2$ :

$$\begin{aligned} \int_0^1 1 \cdot \arctan x \, dx &= x \arctan x \Big|_0^1 - \int_0^1 x \frac{1}{1+x^2} \, dx \\ &= \frac{\pi}{4} - \int_1^2 \frac{1}{u} \left( \frac{1}{2} du \right) = \frac{\pi}{4} - \frac{1}{2} \ln u \Big|_1^2 \\ &= \frac{\pi}{4} - \frac{1}{2} \ln 2 \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad \int_1^2 \frac{x^2 - 2x - 4}{x^3 + 2x^2} \, dx &= \int_1^2 \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x+2} \, dx & \begin{aligned} x^2 - 2x - 4 \\ &= A(x^2 + 2x) + B(x + 2) \\ &\quad + Cx^2 \\ &= (A + C)x^2 + (2A + B)x \\ &\quad + 2B \end{aligned} \\ &= \int_1^2 -\frac{2}{x^2} + \frac{1}{x+2} \, dx & \begin{aligned} 2B &= -4 \longrightarrow B = -2 \\ 2A + B &= -2 \longrightarrow A = 0 \\ A + C &= 1 \longrightarrow C = 1 \end{aligned} \\ &= \left( \frac{2}{x} + \ln |x+2| \right) \Big|_1^2 \\ &= -1 + \ln 4 - \ln 3 \\ &= -1 + \ln \frac{4}{3} \end{aligned}$$

5. (a) Evaluate:  $\int (\tan x + \sin^2 x \cos x) \sin 2x \, dx$   
 [Suggestion: Use the double angle formula for  $\sin 2x$ .]
- (b) Evaluate:  $\int \sqrt{1 - 4x^2} \, dx$

—————SOLUTION—————

(a) Let  $u = \cos x$ :

$$\begin{aligned} & \int \left( \frac{\sin x}{\cos x} + \sin^2 x \cos x \right) 2 \sin x \cos x \, dx \\ &= \int 2 \sin^2 x + 2 \sin^3 x \cos^2 x \, dx \\ &= \int 1 - \cos 2x \, dx + \int 2(1 - \cos^2 x) \cos^2 x \sin x \, dx \\ &= x - \frac{1}{2} \sin 2x + \int 2(u^4 - u^2) \, du \\ &= x - \frac{1}{2} \sin 2x + \frac{2 \cos^5 x}{5} - \frac{2 \cos^3 x}{3} + C \end{aligned}$$

$\begin{aligned} & \text{(b) } \int \sqrt{1 - 4x^2} \, dx \\ &= \int \cos \theta \left( \frac{1}{2} \cos \theta \right) d\theta \\ &= \frac{1}{4} \int 1 + \cos 2\theta \, d\theta \\ &= \frac{1}{4} \left( \theta + \frac{1}{2} \sin 2\theta \right) \\ &= \frac{1}{4} \arcsin 2x + \frac{1}{4} \sin \theta \cos \theta \\ &= \frac{1}{4} \arcsin 2x + \frac{1}{4} (2x) \sqrt{1 - 4x^2} \\ &= \frac{1}{4} \arcsin 2x + \frac{1}{2} x \sqrt{1 - 4x^2} + C \end{aligned}$	$\begin{aligned} 2x &= \sin \theta \\ x &= \frac{1}{2} \sin \theta \\ \arcsin 2x &= \theta \\ dx &= \frac{1}{2} \cos \theta \, d\theta \end{aligned}$
--	---

6. (a) Which of the following improper integrals is/are convergent? Show why.

$$(i) \int_0^1 \frac{\ln(x+2)}{x\sqrt{x}} dx \quad (ii) \int_1^\infty x e^{-2x} dx$$

$$(iii) \int_1^\infty \frac{x \arctan x}{\sqrt[3]{x^7+7}} dx$$

$$(b) \text{ Find } \lim_{x \rightarrow \infty} e^{\frac{x+\ln x}{\sqrt{x^2+x+1}}}.$$

—————SOLUTION—————

$$(a, i) \int_0^1 \frac{\ln(x+2)}{x\sqrt{x}} dx \geq \int_0^1 \frac{\ln 2}{x^{3/2}} dx = \infty. \text{ Div}$$

$$\begin{aligned} (ii) \int_1^\infty x e^{-2x} dx &= -\frac{1}{2} x e^{-2x} \Big|_1^\infty - \int_1^\infty -\frac{1}{2} e^{-2x} dx \\ &= -\frac{1}{2} \frac{x}{e^{2x}} \Big|_1^\infty - \frac{1}{4} e^{-2x} \Big|_1^\infty \\ &= \frac{1}{2e^2} + \frac{1}{4e^2} = \frac{3}{4e^2}. \text{ Convergent} \end{aligned}$$

$$(iii) \int_1^\infty \frac{x \arctan x}{\sqrt[3]{x^7+7}} dx \leq \int_1^\infty \frac{\pi/2}{\sqrt[3]{x^7+7}} dx,$$

and the second integral is convergent by the comparison with

$$\int_1^\infty \frac{1}{x^{7/3}} dx.$$

$$(b) \lim_{x \rightarrow \infty} e^{\frac{x+\ln x}{\sqrt{x^2+x+1}}} = \lim_{x \rightarrow \infty} e^{\frac{x}{\sqrt{x^2+x+1}} + \frac{\ln x}{\sqrt{x^2+x+1}}} = e^{1+0} = e.$$

7. State, for each series, whether it converges absolutely, converges conditionally or diverges. Name a test which supports each conclusion and show the work to apply the test.

$$\begin{array}{ll} \text{(a)} \sum_{n=1}^{\infty} (-1)^n \left(1 + \frac{1}{n} + \frac{1}{n^2}\right)^n & \text{(b)} \sum_{n=0}^{\infty} \frac{n^2 3^n \ln(n+2)}{2^{2n+2}} \\ \text{(c)} \sum_{n=0}^{\infty} \frac{(-1)^n (n^2 + 3)}{n^3 + 4} & \end{array}$$

—————SOLUTION—————

For each series,  $a_n$  will denote the  $n$ th term of the series.

(a) Since  $a_n \geq 1$  for all  $n$ ,  $\lim a_n \neq 0$  and the series diverges by the Test for Divergence.

(b) By L'Hopital's Rule, as  $n$  and  $x$  go to infinity,

$$\lim \frac{\ln(n+3)}{\ln(n+2)} = \lim \frac{\ln(x+3)}{\ln(x+2)} = \lim \frac{1/(x+3)}{1/(x+2)} \longrightarrow 1. \text{ So}$$

$$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)^2}{n^2} \frac{\ln(n+3)}{\ln(n+2)} \frac{3^{n+1}}{3^n} \frac{2^{2n+2}}{2^{2n+4}} \longrightarrow 1 \cdot 1 \cdot 3 \cdot \frac{1}{2^2} = \frac{3}{4} < 1.$$

Thus, the series converges absolutely by the Ratio Test.

(c) Clearly  $a_n$  is alternating and converges to zero.

Since  $\left( \frac{x^2 + 3}{x^3 + 4} \right)' = -\frac{x(x^2 + 3x + 2)}{(x^3 + 4)^2} < 0$  for  $x > 0$ , the terms

form a decreasing sequence, and the Alternating Series Test says the series is convergent. The series  $\sum |a_n|$  is divergent by the Limit Comparison Test with  $b_n = \frac{1}{n}$ . Thus,  $\sum a_n$  converges conditionally.

8. (a) Find the interval of convergence of the power series

$$\sum_{n=0}^{\infty} \frac{n(x+1)^n}{5^n \sqrt{n^2+4}}.$$

Remember to check the endpoints if applicable.

(b) Sketch the graph of the polar equation  $r = 3 + 2 \sin \theta$ , and find the area which is both inside the graph and above the  $x$ -axis.

### —————SOLUTION—————

(a) The center of the series is  $x_0 = -1$ , and the  $n$ th coefficient is  $c_n = \frac{1}{5^n \sqrt{n^2+4}}$ .

$$\left| \frac{c_{n+1}}{c_n} \right| = \frac{5^n \sqrt{n^2+4}}{5^{n+1} \sqrt{(n+1)^2+4}} \longrightarrow \frac{1}{5},$$

so the radius of convergence is  $R = 5$ , and the series converges for  $x$  in the open interval  $(-1-5, -1+5) = (-6, 4)$ .

At  $x = -6$  the series is  $\sum \frac{n(-1)^n}{\sqrt{n^2+4}}$  which is divergent.

At  $x = 4$  the series is  $\sum \frac{n}{\sqrt{n^2+4}}$  which is divergent.


The interval of convergence is  $(-6, 4)$ .

$$(b) \quad r(0) = 3$$

$$r(\pi/2) = 5$$

$$r(\pi) = 3$$

$$r(3\pi/2) = 1$$



$$\begin{aligned} A &= \frac{1}{2} \int_0^{\pi} (3 + 2 \sin \theta)^2 d\theta \\ &= \frac{1}{2} \int_0^{\pi} 9 + 12 \sin \theta + 4 \sin^2 \theta d\theta \\ &= \frac{9\pi}{2} - 6 \cos \theta \Big|_0^{\pi} + \int_0^{\pi} 1 - \cos 2\theta d\theta \\ &= \frac{9\pi}{2} + 12 + \pi = \frac{11\pi}{2} + 12. \end{aligned}$$



9. (a) Find the first four nonzero terms of the Maclaurin series (i.e., power series centered at 0) for the function

$$g(x) = e^{-x^2}.$$

- (b) Find the sum of the first four terms of the Maclaurin series for the derivative of the function  $f(x) = \sum_{n=0}^{\infty} \frac{x^n}{(n+1)^3}$ .

- (c) Use the answer in part (b) to approximate  $f'(-\frac{1}{2})$  with an error of less than .01.

—————SOLUTION—————

- (a) Replace  $x$  by  $-x^2$  in the series for  $e^x$ :

$$\begin{aligned} e^x &= \sum_{n=0}^{\infty} \frac{x^n}{n!} \\ e^{-x^2} &= \sum_{n=0}^{\infty} \frac{(-x^2)^n}{n!} \\ &= \sum_{n=0}^{\infty} \frac{(-)^n x^{2n}}{n!} = 1 - x^2 + \frac{x^4}{2!} + \frac{x^6}{3!} \pm \dots \end{aligned}$$

$$(b) \quad f(x) = \sum_{n=0}^{\infty} \frac{x^n}{(n+1)^3} = 1 + \frac{x}{2^3} + \frac{x^2}{3^3} + \frac{x^3}{4^3} + \frac{x^4}{5^3} + \dots$$

$$f'(x) = \sum_{n=1}^{\infty} \frac{nx^{n-1}}{(n+1)^3} = \frac{1}{2^3} + \frac{2x}{3^3} + \frac{3x^2}{4^3} + \frac{4x^3}{5^3} + \dots$$

$$f'(-\frac{1}{2}) = \boxed{\frac{1}{2^3} - \frac{2}{3^3 2} + \frac{3}{4^3 2^2}} + \frac{4}{5^3 2^3} + \dots$$

The boxed sum has error not more than  $\frac{4}{5^3 2^3} = \frac{1}{250} < .01$ .

10. (a) Graph  $4x^2 + 36y^2 + 9z^2 + 16x - 20 = 0$ , and graph the trace of the answer to (a) in the  $xy$ -plane.

(b) Sketch the portion of the graph of  $y = 1 - x^2$  which is in the first octant.

—————SOLUTION—————

$$(a) \quad 4x^2 + 16x + 36y^2 + 9z^2 - 20 = 0$$



$$4[x^2 + 4x] + 36y^2 + 9z^2 - 20 = 0$$

$$4[(x + 2)^2 - 4] + 36y^2 + 9z^2 - 20 = 0$$

$$4(x + 2)^2 - 16 + 36y^2 + 9z^2 - 20 = 0$$

$$4(x + 2)^2 + 36y^2 + 9z^2 = 36$$

$$\frac{(x + 2)^2}{9} + y^2 + \frac{z^2}{4} = 1$$



(b) The graph is the cylinder obtained by moving the graph of  $y = 1 - x^2$ ,  $x \geq 0$  in the  $z$ -direction.

